arXiv:math/0510509v1 [math.GR] 24 Oct 2005

ON FIXED-POINT SETS IN THE BOUNDARY OF A CAT(0) SPACE

TETSUYA HOSAKA

Abstract. In this paper, we investigate the fixed-point set of an element of a CAT(0) group in its boundary. Suppose that a group G acts geometrically on a CAT(0) space X. Let $g \in G$ and let $\mathcal{F}_g$ be the fixed-point set of g in the boundary ∂X. Then we show that $\mathcal{F}_g = L(Z_g)$, where Z_g is the centralizer of g (i.e. $Z_g = \{v \in G \,|\, gv = vg\}$) and $L(Z_g)$ is the limit set of Z_g in ∂X. Thus we obtain that $\mathcal{F}_g \neq \emptyset$ if and only if the set Z_g is infinite. We also show that if g is a hyperbolic isometry, then $\mathcal{F}_g = \partial \operatorname{Min}(g)$, where $\partial \operatorname{Min}(g)$ is the boundary of the minimal set $\operatorname{Min}(g)$ of g. This implies that the fixed-point set $\mathcal{F}_g$ and the periodic-point set $\mathcal{P}_g$ of g in ∂X have suspension forms.

1. Introduction and preliminaries

The purpose of this paper is to study the fixed-point set of an element of a CAT(0) group in its boundary.

We say that a metric space (X, d) is a *geodesic space* if for each $x, y \in X$, there exists an isometric embedding $\xi : [0, d(x, y)] \to X$ such that $\xi(0) = x$ and $\xi(d(x, y)) = y$ (such ξ is called a *geodesic*). Also a metric space (X, d) is said to be *proper* if every closed metric ball is compact.

Let (X, d) be a geodesic space and let T be a geodesic triangle in X. A *comparison triangle* for T is a geodesic triangle $\overline{T}$ in the Euclidean plane $\mathbb{R}^2$ with same edge lengths as T. Choose two points x and y in T. Let $\bar{x}$ and $\bar{y}$ denote the corresponding points in $\overline{T}$. Then the inequality

$$d(x, y) \leq d_{\mathbb{R}^2}(\bar{x}, \bar{y})$$

Date: May 2, 2005.

2000 *Mathematics Subject Classification.* 57M07.

Key words and phrases. boundaries of CAT(0) groups.

Partly supported by the Grant-in-Aid for Young Scientists (B), The Ministry of Education, Culture, Sports, Science and Technology, Japan. (No. 15740029).

is called the *CAT(0)-inequality*, where $d_{\mathbb{R}^2}$ is the natural metric on $\mathbb{R}^2$. A geodesic space (X, d) is called a *CAT(0) space* if the CAT(0)-inequality holds for all geodesic triangles T and for all choices of two points x and y in T.

Let X be a proper CAT(0) space and $x_0 \in X$. The *boundary of X with respect to x_0*, denoted by $\partial_{x_0} X$, is defined as the set of all geodesic rays issuing from x_0. Then we define a topology on $X \cup \partial_{x_0} X$ by the following conditions:

(1) X is an open subspace of $X \cup \partial_{x_0} X$.

(2) For $\alpha \in \partial_{x_0} X$ and $r, \epsilon > 0$, let

$$U_{x_0}(\alpha; r, \epsilon) = \{x \in X \cup \partial_{x_0} X \mid x \notin B(x_0, r),\ d(\alpha(r), \xi_x(r)) < \epsilon\},$$

where $\xi_x : [0, d(x_0, x)] \to X$ is the geodesic from x_0 to x ($\xi_x = x$ if $x \in \partial_{x_0} X$). Then for each $\epsilon_0 > 0$, the set

$$\{U_{x_0}(\alpha; r, \epsilon_0) \mid r > 0\}$$

is a neighborhood basis for α in $X \cup \partial_{x_0} X$.

This is called the *cone topology* on $X \cup \partial_{x_0} X$. It is known that $X \cup \partial_{x_0} X$ is a metrizable compactification of X ([1], [2]).

Let X be a geodesic space. Two geodesic rays $\xi, \zeta : [0, \infty) \to X$ are said to be *asymptotic* if there exists a constant N such that $d(\xi(t), \zeta(t)) \leq N$ for each $t \geq 0$. It is known that for each geodesic ray ξ in X and each point $x \in X$, there exists a unique geodesic ray ξ' issuing from x such that ξ and ξ' are asymptotic.

Let x_0 and x_1 be two points of a proper CAT(0) space X. Then there exists a unique bijection $\Phi : \partial_{x_0} X \to \partial_{x_1} X$ such that ξ and $\Phi(\xi)$ are asymptotic for each $\xi \in \partial_{x_0} X$. It is known that $\Phi : \partial_{x_0} X \to \partial_{x_1} X$ is a homeomorphism ([1], [2]).

Let X be a proper CAT(0) space. The asymptotic relation is an equivalence relation in the set of all geodesic rays in X. The *boundary of X*, denoted by ∂X, is defined as the set of asymptotic equivalence classes of geodesic rays. The equivalence class of a geodesic ray ξ is denoted by $\xi(\infty)$. For each $x_0 \in X$ and each $\alpha \in \partial X$, there exists a unique element $\xi \in \partial_{x_0} X$ with $\xi(\infty) = \alpha$. Thus we may identify ∂X with $\partial_{x_0} X$ for each $x_0 \in X$.

Let X be a proper CAT(0) space and G a group which acts on X by isometries. For each element $g \in G$ and each geodesic ray $\xi : [0, \infty) \to X$, a map $g\xi : [0, \infty) \to X$ defined by $(g\xi)(t) := g(\xi(t))$ is also a geodesic ray. If geodesic rays ξ and ξ' are asymptotic, then $g\xi$ and $g\xi'$ are also asymptotic. Thus g induces a homeomorphism of ∂X and G acts on ∂X.

A *geometric* action on a CAT(0) space is an action by isometries which is proper ([1, p.131]) and cocompact. We note that every CAT(0) space on which a group acts geometrically is a proper space ([1, p.132]).

Details of CAT(0) spaces and their boundaries are found in [1] and [2].

Suppose that a group G acts geometrically on a CAT(0) space X. For a subset $A \subset G$, the limit set $L(A)$ of A is defined as $L(A) = \overline{Ax_0} \cap \partial X$, where $x_0 \in X$ and $\overline{Ax_0}$ is the closure of the orbit Ax_0 in $X \cup \partial X$. We note that the limit set $L(A)$ is determined by A and not depend on the point x_0. For $g \in G$, we define F_g and $\mathcal{F}_g$ as the fixed-point sets of g in X and ∂X respectively, that is, $F_g = \{x \in X \mid gx = x\}$ and $\mathcal{F}_g = \{\alpha \in \partial X \mid g\alpha = \alpha\}$. By [1, Corollary II.2.8 (1)], we see that $F_g \neq \emptyset$ if and only if the order $o(g)$ of g is finite, since the action of G on X is proper. In this paper, we investigate $\mathcal{F}_g$. We prove the following theorem in Section 2.

Theorem 1.1. *Suppose that a group G acts geometrically on a CAT(0) space X. For $g \in G$, $\mathcal{F}_g = L(Z_g)$, where Z_g is the centralizer of g, i.e., $Z_g = \{v \in G \mid gv = vg\}$.*

For a subset $A \subset G$, $L(A) \neq \emptyset$ if and only if A is infinite. Hence Theorem 1.1 implies the following corollary.

Corollary 1.2. *Suppose that a group G acts geometrically on a CAT(0) space X. For $g \in G$, $\mathcal{F}_g \neq \emptyset$ if and only if Z_g is infinite.*

We can obtain the following corollary from Corollary 1.2.

Corollary 1.3. *Suppose that a group G acts geometrically on CAT(0) spaces X and X'. For $g \in G$, $\mathcal{F}_g \neq \emptyset$ if and only if $\mathcal{F}'_g \neq \emptyset$, where $\mathcal{F}_g$ and $\mathcal{F}'_g$ are the fixed-point sets of g in ∂X and $\partial X'$ respectively.*

Let X be a CAT(0) space and let g be an isometry of X. The *translation length* of g is the number $|g| := \inf\{d(x, gx) \mid x \in X\}$, and the *minimal set* of g is defined as $\mathrm{Min}(g) := \{x \in X \mid d(x, gx) = |g|\}$. An isometry g of X is said to be *hyperbolic*, if $\mathrm{Min}(g) \neq \emptyset$ and $|g| > 0$ (cf. [1]).

We also prove the following theorem in Section 2.

Theorem 1.4. *Let g be a hyperbolic isometry of a proper CAT(0) space X. Then the fixed-point set of g in ∂X is $\mathcal{F}_g = \partial \mathrm{Min}(g)$, where $\partial \mathrm{Min}(g)$ is the boundary of the minimal set $\mathrm{Min}(g)$ of g.*

For a hyperbolic isometry g of a CAT(0) space X, $\mathrm{Min}(g)$ is the union of the axes of g and $\mathrm{Min}(g)$ splits as a product $Y_1 \times \mathrm{Im}\,\sigma$, where $\sigma : \mathbb{R} \to X$ is an axis of g ([1, p.231]). Hence $\partial \mathrm{Min}(g)$ is the suspension

$\partial Y_1 * \{g^{-\infty}, g^{\infty}\}$. Theorem 1.4 implies that the fixed-point set $\mathcal{F}_g$ has a suspension form. Let $\mathcal{P}_g$ be the periodic-point set of g in ∂X. Then

$$\mathcal{P}_g = \bigcup_{n\in\mathbb{N}} \mathcal{F}_{g^n} = \bigcup_{n\in\mathbb{N}} \partial \operatorname{Min}(g^n).$$

Hence the periodic-point set $\mathcal{P}_g$ of g is also a suspension form.

We obtain the following corollary from Theorems 1.1 and 1.4.

Corollary 1.5. *Suppose that a group G acts geometrically on a CAT(0) space X. For $g \in G$ such that $o(g) = \infty$, $\mathcal{F}_g = L(Z_g) = \partial \operatorname{Min}(g)$.*

2. Proof of the main theorems

We first prove Theorem 1.1.

Proof of Theorem 1.1. Suppose that a group G acts geometrically on a CAT(0) space X. Let $g \in G$ and let $x_0 \in X$.

We first show that $\mathcal{F}_g \subset L(Z_g)$. Let $\alpha \in \mathcal{F}_g$ and let $\xi : [0, \infty) \to X$ be the geodesic ray such that $\xi(0) = x_0$ and $\xi(\infty) = \alpha$. Since $\alpha \in \mathcal{F}_g$, $g\alpha = \alpha$. Hence the geodesic rays ξ and $g\xi$ are asymptotic, and there exists a number $M > 0$ such that $d(\xi(t), g\xi(t)) \leq M$ for any $t \geq 0$. Since the action of G on X is cocompact and X is proper, $GB(x_0, N) = X$ for some $N > 0$. For each $i \in \mathbb{N}$, there exists $v_i \in G$ such that $d(\xi(i), v_i x_0) \leq N$. Then

$$\begin{aligned} d(x_0, v_i^{-1} g v_i x_0) &= d(v_i x_0, g v_i x_0) \\ &\leq d(v_i x_0, \xi(i)) + d(\xi(i), g\xi(i)) + d(g\xi(i), g v_i x_0) \\ &\leq 2N + M, \end{aligned}$$

because $d(v_i x_0, \xi(i)) \leq N$ and $d(\xi(i), g\xi(i)) \leq M$. Since the action of G on X is proper, the set $\{h \in G |\, d(x_0, hx_0) \leq 2N + M\}$ is finite. Hence there exists $g' \in G$ such that $\{i \in \mathbb{N} |\, v_i^{-1} g v_i = g'\}$ is infinite. Let $\{i_j |\, j \in \mathbb{N}\} = \{i \in \mathbb{N} |\, v_i^{-1} g v_i = g'\}$. Then $v_{i_j}^{-1} g v_{i_j} = g'$ for each $j \in \mathbb{N}$, and

$$v_{i_j}^{-1} g v_{i_j} = g' = v_{i_1}^{-1} g v_{i_1}.$$

Hence $g = (v_{i_j} v_{i_1}^{-1}) g (v_{i_1} v_{i_j}^{-1})$. Thus $v_{i_j} v_{i_1} \in Z_g$ for each $j \in \mathbb{N}$. Here we note that $v_{i_j} v_{i_1}^{-1} \neq v_{i_{j'}} v_{i_1}^{-1}$ if $j \neq j'$. Since $\{v_{i_j} x_0 |\, j \in \mathbb{N}\}$ converges to α in $X \cup \partial X$, $\{v_{i_j} v_{i_1}^{-1} x_0 |\, j \in \mathbb{N}\}$ converges to α. Hence $\alpha \in L(\{v_{i_j} v_{i_1}^{-1} |\, j \in \mathbb{N}\}) \subset L(Z_g)$. Thus $\mathcal{F}_g \subset L(Z_g)$.

Next, we show that $\mathcal{F}_g \supset L(Z_g)$. Let $\alpha \in L(Z_g)$. There exists a sequence $\{v_i |\, i \in \mathbb{N}\} \subset Z_g$ such that $\{v_i x_0 |\, i \in \mathbb{N}\}$ converges to α in $X \cup \partial X$. For each $i \in \mathbb{N}$, $g v_i = v_i g$. Here the sequence $\{g v_i x_0 |\, i \in \mathbb{N}\}$ converges to $g\alpha$ and $\{v_i g x_0 |\, i \in \mathbb{N}\}$ converges to α. Thus $g\alpha = \alpha$, i.e., $\alpha \in \mathcal{F}_g$.

Therefore $\mathcal{F}_g = L(Z_g)$. $\square$

Next we prove Theorem 1.4.

Proof of Theorem 1.4. Let g be a hyperbolic isometry of a proper CAT(0) space X and let $x_0 \in \operatorname{Min}(g)$.

We first show that $\mathcal{F}_g \subset \partial \operatorname{Min}(g)$. Let $\alpha \in \mathcal{F}_g$ and let $\xi : [0, \infty) \to X$ be the geodesic ray such that $\xi(0) = x_0$ and $\xi(\infty) = \alpha$. Since $\alpha \in \mathcal{F}_g$, $g\alpha = \alpha$. Hence the geodesic rays ξ and $g\xi$ are asymptotic. Now $d(x_0, gx_0) = |g|$ because $x_0 \in \operatorname{Min}(g)$. Hence

$$d(\xi(t), g\xi(t)) \leq d(\xi(0), g\xi(0)) = d(x_0, gx_0) = |g|.$$

This means that $d(\xi(t), g\xi(t)) = |g|$ and $\xi(t) \in \operatorname{Min}(g)$ for each $t \geq 0$. Hence $\operatorname{Im} \xi \subset \operatorname{Min}(g)$ and $\alpha \in \partial \operatorname{Min}(g)$. Thus $\mathcal{F}_g \subset \partial \operatorname{Min}(g)$.

Next we show that $\mathcal{F}_g \supset \partial \operatorname{Min}(g)$. Let $\alpha \in \partial \operatorname{Min}(g)$ and let $\xi : [0, \infty) \to X$ be the geodesic ray such that $\xi(0) = x_0$ and $\xi(\infty) = \alpha$. Then $\operatorname{Im} \xi \subset \operatorname{Min}(g)$, since $\alpha \in \partial \operatorname{Min}(g)$ and $\operatorname{Min}(g)$ is convex. Hence $\xi(t) \in \operatorname{Min}(g)$ and $d(\xi(t), g\xi(t)) = |g|$ for any $t \geq 0$. This means that ξ and $g\xi$ are asymptotic and $\alpha = g\alpha$, i.e., $\alpha \in \mathcal{F}_g$. Thus $\mathcal{F}_g \supset \partial \operatorname{Min}(g)$.

Therefore $\mathcal{F}_g = \partial \operatorname{Min}(g)$. $\square$

3. REMARKS

Let g be a hyperbolic isometry of a proper CAT(0) space X. For each $n \in \mathbb{N}$, $\mathcal{F}_{g^n} = \partial \operatorname{Min}(g^n)$ by Theorem 1.4. We note that an axis of g is also an axis of g^n for each $n \in \mathbb{N}$. Then the minimal set $\operatorname{Min}(g^n)$ of g^n splits as a product $\operatorname{Min}(g^n) = Y_n \times \operatorname{Im} \sigma$, where $\sigma : \mathbb{R} \to X$ is an axis of g (cf. [1]). Here $Y_n \subset Y_{kn}$ for each $n, k \in \mathbb{N}$. The fixed-point set of g in ∂X is

$$\mathcal{F}_g = \partial \operatorname{Min}(g) = \partial(Y_1 \times \operatorname{Im} \sigma) = \partial Y_1 * \{g^{\infty}, g^{-\infty}\}.$$

Also the periodic-point set of g is

$$\begin{aligned} \mathcal{P}_g &= \bigcup_{n \in \mathbb{N}} \mathcal{F}_{g^n} = \bigcup_{n \in \mathbb{N}} \partial \operatorname{Min}(g^n) \\ &= \bigcup_{n \in \mathbb{N}} \partial(Y_n \times \operatorname{Im} \sigma) = (\bigcup_{n \in \mathbb{N}} \partial Y_n) * \{g^{\infty}, g^{-\infty}\}. \end{aligned}$$

Thus the fixed-point set and the periodic-point set of g have suspension forms.

Let A be the union of geodesic lines which are parallel to an axis σ of g. Then A splits as a product $Y \times \operatorname{Im} \sigma$. By the above argument, $\mathcal{P}_g = \bigcup_{n \in \mathbb{N}} \partial \operatorname{Min}(g^n) \subset \partial A$ and $\bigcup_{n \in \mathbb{N}} \partial Y_n \subset \partial Y$.

Here the following problem arises.

Problem 3.1. Is it always the case that $\mathcal{P}_g = \partial A$ (i.e. $\bigcup_{n\in\mathbb{N}} \partial Y_n = \partial Y$)?

Let g be a hyperbolic isometry of a Gromov hyperbolic space X. By an easy argument, we see that the fixed-point set $\mathcal{F}_g$ of g in the boundary ∂X is the two-points set $\{g^{\infty}, g^{-\infty}\}$. Also for each $\alpha \in \partial X \setminus \{g^{-\infty}\}$, the sequence $\{g^i\alpha \,|\, i \in \mathbb{N}\}$ converges to g^{∞} in ∂X.

Here the following problem arises.

Problem 3.2. Let g be a hyperbolic isometry of a proper CAT(0) space X and let A be the union of geodesic lines which are parallel to an axis σ of g. Is it the case that for each $\alpha \in \partial X \setminus \partial A$, the sequence $\{g^i\alpha \,|\, i \in \mathbb{N}\}$ converges to g^{∞} in ∂X?

Also the following problem arises.

Problem 3.3. Suppose that a group G acts geometrically on a CAT(0) space X. Is it always the case that there do not exist $g \in G$ and $\alpha, \beta \in \partial X$ such that $\limsup_{n\to\infty} d_{\partial X}(g^n\alpha, g^n\beta) > 0$ and $\liminf_{n\to\infty} d_{\partial X}(g^n\alpha, g^n\beta) = 0$?

REFERENCES

[1] M.R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Springer-Verlag, Berlin, 1999.

[2] E. Ghys and P. de la Harpe (ed), *Sur les Groups Hyperboliques d'apres Mikhael Gromov*, Progr. Math. vol. 83, Birkhäuser, Boston MA, 1990.

[3] M. Gromov, *Hyperbolic groups*, Essays in group theory (S. M. Gersten, ed.), M.S.R.I. Publ. 8, 1987, pp. 75-264.

DEPARTMENT OF MATHEMATICS, UTSUNOMIYA UNIVERSITY, UTSUNOMIYA, 321-8505, JAPAN

E-mail address: `hosaka@cc.utsunomiya-u.ac.jp`